Section 1: Pain Intensity
The Pain comes and goes and is vary mild.
The pain is mild and does not vary much.
The pain is moderate and does not vary much.
The pain comes and goes and is very severe.
The pain is severe and does not vary much.
Section 2: Personal Care (washing, Dressing, etc.)
I would not have to change the way of washing or dressing in order to avoid pain.
I do not normally change my way of washing and dressing even though it causes some pain.
Washing and dressing increases the pain, but I manage not to change my way of doing it.
Washing and dressing increases the pain and I find it necessary to change my way of doing it.
Because of the pain, I am unable to do some washing and dressing without help.
Because of the pain, I am unable to do any washing and dressing without help.
Section 3: Lifting
I can lift heavy weights without extra pain.
I can lift heavy weights but it gives me extra pain.
Pain prevents me from lifting heavy weights off the floor but I can if they are in convenient places.
Pain prevents me from lifting heavy weights off the floor.
Pain prevents me from lifting heavy weights, but I can manage medium weights conveniently positioned.
I can only lift very light weights at the most

Section 4: Walking	
I have no pain walking.	
I have some pain on walking, but it does not increase with distance.	
I cannot walk more than one mile without increasing pain.	
I cannot walk more than ½ mile without increasing pain.	
I cannot walk more than ¼ mile without increasing pain.	
I cannot walk at all without increasing pain.	
Section 5: Sitting	
I can it in any chair as long as I like.	
I only sit in my favorite chair as long as I like.	
Pain prevents me from sitting more than 1 hour.	
Pain prevents me from sitting more than ½ hour.	
Pain prevents me from sitting more than 10 minutes	
I avoid sitting because it increases my pain right away.	
Section 6: Standing	
I can stand as long as I want without extra pain.	
I have some pain on standing, but it does not increase with time.	
I cannot stand for longer than 1 hour without increasing pain	
I cannot stand for longer than ½ hour without increasing pain.	
I cannot stand for longer than 10 minutes without increasing pain.	
I avoid standing because it increases the pain right away.	

Section 7: Sleeping
I get pain in bed, but it does not prevent me from sleeping well.
Because of pain, my normal night's sleep is reduced by less than 1/4.
Because of pain, my normal night's sleep is reduced by less than ½.
Because of pain, my normal night's sleep is reduced by less than 3/4.
Pain prevents me from sleeping at all.
Pain prevents me from sleeping at all.
Section 8: Social Life
My social life is normal and gives me no extra pain.
My social life is normal but it increases the degree of pain
Pain has no significant effect on my social life apart from limiting my more energetic interests,
Pain has restricted my social life and I do not go out as often.
Pain has restricted my social to my home
Section 9: Traveling
I get no pain while traveling
I get some pain while traveling, but none of my usual forms of travel makes it any worse.
I get extra pain while traveling, but it does not compel me to seek alternative form of travel.
I get extra pain while traveling, which compels me to seek alternative forms of travel.
Pain restricts all forms of travel.
Pain prevents all forms of travel except that down by lying down

#### **Section 10: Changing Degree of Pain**

My pain is rapidly getting better.
My Pain fluctuates, but is definitely getting better.
My pain seem to be getting better, but improvement is slow at present.
Pain is neither getting better nor worse.
Pain is gradually worsening.
Pain is rapidly worsening.